





Plant respiration: Controlled by photosynthesis or biomass?

Alessio Collalti^{1,2}  | Mark G. Tjoelker³  | Günter Hoch⁴  | Annikki Mäkelä⁵  |
Gabriele Guidolotti⁶  | Mary Heskell⁷  | Gai Petit⁸  | Michael G. Ryan^{9,10}  |
Giovanna Battipaglia¹¹  | Giorgio Matteucci¹  | Iain Colin Prentice^{12,13,14} 

¹Institute for Agriculture and Forestry Systems in the Mediterranean, National Research Council of Italy (CNR-ISAFOM), Rende (CS), Italy

²Department of Innovation in Biological, Agro-food and Forest Systems, University of Tuscia, Viterbo, Italy

³Hawkesbury Institute for the Environment, Western Sydney University, Penrith, NSW, Australia

⁴Department of Environmental Sciences – Botany, University of Basel, Basel, Switzerland

⁵Institute for Atmospheric and Earth System Research (INAR), Faculty of Science and Faculty of Agriculture and Forestry, University of Helsinki, Helsinki, Finland

⁶Institute of Research on Terrestrial Ecosystem, National Research Council of Italy (CNR-IRET), Rome, Italy

⁷Department of Biology, Macalester College, Saint Paul, MN, USA

⁸Department of Land, Environment, Agriculture and Forestry, University of Padova, Padova, Italy

⁹Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, CO, USA

¹⁰USDA Forest Service, Rocky Mountain Experiment Station, Fort Collins, CO, USA

¹¹Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Campania "L. Vanvitelli", Caserta, Italy

¹²AXA Chair of Biosphere and Climate Impacts, Department of Life Sciences, Imperial College London, Ascot, UK

¹³Department of Biological Sciences, Macquarie University, North Ryde, NSW, Australia

¹⁴Department of Earth System Science, Ministry of Education Key Laboratory for Earth System Modeling, Tsinghua University, Beijing, China

Correspondence

Alessio Collalti, Institute for Agriculture and Forestry Systems in the Mediterranean, National Research Council of Italy (CNR-ISAFOM), 87036 Cosenza, Italy.
Email: alessio.collalti@cnr.it

Funding information

European Research Council, Grant/Award Number: 787203

Abstract

Two simplifying hypotheses have been proposed for whole-plant respiration. One links respiration to photosynthesis; the other to biomass. Using a first-principles carbon balance model with a prescribed live woody biomass turnover, applied at a forest research site where multidecadal measurements are available for comparison, we show that if turnover is fast the accumulation of respiring biomass is low and respiration depends primarily on photosynthesis; while if turnover is slow the accumulation of respiring biomass is high and respiration depends primarily on biomass. But the first scenario is inconsistent with evidence for substantial carry-over of fixed carbon between years, while the second implies far too great an increase in respiration during stand development—leading to depleted carbohydrate reserves and an unrealistically high mortality risk. These two mutually incompatible hypotheses are thus both incorrect. Respiration is *not* linearly related either to photosynthesis or to biomass, but it is more strongly controlled by recent photosynthates (and reserve availability) than by total biomass.

KEYWORDS

biomass accumulation, carbon use efficiency, gross primary production, maintenance respiration, metabolic scaling theory, net primary production, nonstructural carbohydrates, plant respiration

1 | INTRODUCTION

The amount of carbon that accumulates in actively growing stands of vegetation depends on the balance of photosynthesis (gross primary production, P) and whole-plant (autotrophic) respiration (R). The difference between these fluxes is the net primary production (P_n). Most annual P_n is allocated to structural growth (G), but some is stored as nonstructural carbohydrates (NSC, mostly starch and sugars), some is released back to the atmosphere in the form of biogenic volatile organic compounds (BVOCs), and some is exuded to the rhizosphere (Chapin et al., 2006). The fraction of P that accumulates in biomass, and the fraction that returns to the atmosphere through plant metabolism, are crucial quantities that determine the sign and magnitude of the global climate-carbon feedback—which remains one of the greatest sources of uncertainty in the global carbon cycle (Friedlingstein et al., 2014). But despite many ecophysiological studies aiming to understand P_n and R dynamics during stand development, a general understanding is still lacking.

Some authors have hypothesized a constant $P_n:P$ (carbon use efficiency, equivalent to $1 - (R:P)$ ratio, with R tightly constrained by P irrespective of biomass, climate, tree species and stand age (e.g. Gifford, 2003; Van Oijen, Schapendonk, & Hoglind, 2010). Waring, Landsberg, and Williams (1998, W98 hereafter) indicated a universal $P_n:P$ of ~0.5. Since, ultimately, R depends on the matter produced by photosynthesis, Gifford (2003) suggested that these two processes must be tightly balanced over the longer term—making R proportional to P , consistent with W98. He argued that prescribing P_n (or R) as a constant fraction of P could be a simpler, and potentially more accurate, alternative to explicit, process-based modelling of R . A number of land vegetation models (reviewed in Collalti & Prentice, 2019) adopt this simplification.

An alternative hypothesis, grounded in metabolic scaling theory, suggests that R should scale with biomass following a power law, $Y = aX^b$ (West, Brown, & Enquist, 1999). According to some studies (e.g. Reich, Tjoelker, Machado, & Oleksyn, 2006, R06 hereafter), R (Y) scales isometrically ($b \sim 1$) with whole-plant carbon (C) or nitrogen (N) contents (X), and this scaling is similar within and among different species, and irrespective of environmental and climatic conditions—which might influence the normalization constant (a), but not the exponent (b). Isometric scaling of R with biomass was assumed in the traditional view of forest dynamics set out for example, by Kira and Shidei (1967) and Odum (1969). In the absence of major disturbances, if R increases in parallel with biomass, then P_n necessarily declines—because ultimately P cannot increase indefinitely, but rather stabilizes at canopy closure. Mori et al. (2010), however, indicated that biomass and R are isometrically related only in young trees, tending towards $b \sim 3/4$ in mature trees. A general value of $3/5$ has also been proposed (Michaletz, Cheng, Kerkhoff, & Enquist, 2014). But however it is interpreted, this scaling hypothesis implies that R depends on biomass, and is related to P only to the extent that P and biomass vary together.

Although many terrestrial vegetation models simulate plant respiration assuming R to be a fixed fraction of P , others more explicitly couple R to biomass and thus only indirectly to P . The most widely used (and observationally supported) mechanistic approach, also adopted here, divides R into growth (R_G) and maintenance (R_M) components (McCree, 1970; Thornley, 1970). R_G is considered to be a fixed fraction of new

tissue growth, independent of temperature, the fraction varying only with the cost of building the compounds constituting the new tissue (Penning de Vries, Brunsting, & van Laar, 1974). Temperature, substrate availability and the demand for respiratory products are considered to control R_M (Cannell & Thornley, 2000). Several studies have investigated the effects of short- and long-term changes in temperature on R_M , mostly at the leaf level (e.g. Heskell et al., 2016; Huntingford et al., 2017). The nature of the temperature responses and the acclimation of R_M are important and much-discussed issues, but they are not considered further here. In contrast, the effects on respiration of woody biomass (the substrate), its accumulation and the transition rate of respiring sapwood into nonrespiring heartwood, have received relatively little attention (Kuptz, Fleischman, Matyssek, & Grams, 2011; Tjoelker, Oleksyn, & Reich, 1999). These latter processes are the focus here.

The fixed-ratio hypothesis of W98 and the scaling hypothesis of R06 could both be used—at least in principle, across the 20 orders of magnitude variation in plant mass—to estimate R and P_n without the need for explicit process-based modelling of R (McMurtrie et al., 2008; Price, Gilooly, Allen, Weitz, & Niklas, 2010). However, they may yield quite different results, and both hypotheses (and their supposed underlying mechanisms) have been subject to criticism (e.g. Agutter & Tuszynski, 2011; Collalti & Prentice, 2019; Collalti et al., 2018, 2019; Keith, Mackey, Berry, Lindenmayer, & Gibbons, 2010; Kozłowski & Konarzewski, 2005; Mäkelä & Valentine, 2001; Medlyn & Dewar, 1999; O'Connor et al., 2007; Price et al., 2012). To our knowledge, there has been no previous attempt to compare these two hypotheses directly, and their consequences for forest carbon balance during stand development, and in the same modelling framework. We attempt to fill this gap by providing illustrative simulations on the long-term trajectories of R , P_n and $P_n:P$, highlighting and discussing the large uncertainty surrounding this issue. The simulations are based on the first principles of mass balance, as adopted in most contemporary vegetation models, and implemented here into a process-based, ecophysiological model that has been tested against detailed time series observations in an intensively monitored research forest site. We show how alternative assumptions about the live woody turnover (live woody biomass is the metabolically active fraction of sapwood: see Supporting Information) map on to the two alternative hypotheses, while seeking an answer to the pivotal question: is R a function of photosynthesis alone (W98's hypothesis), or of biomass alone (R06's hypothesis)? Insight into these conflicting hypotheses on plant respiration would help towards a better mechanistic understanding and correct quantification of the stocks and fluxes that determine the carbon balance of forests.

2 | MATERIALS AND METHODS

2.1 | Theoretical framework

A general equation describing autotrophic respiration (R) is:

$$R = P - P_n = P - (G + G_R), \quad (1)$$

where P and P_n are gross and net primary production, G is structural and litter biomass production and G_R is the flux to NSC reserves

and secondary compounds including, exudates and BVOCs (all in $\text{g C ground area}^{-1} \text{time}^{-1}$). If R is further decomposed into growth (R_G) and maintenance (R_M) respiration (McCree, 1970; Thornley, 1970), then:

$$R = R_G + R_M = g_R G + m_R W_{\text{live}}, \quad (2)$$

where g_R and m_R are the growth and maintenance respiration coefficients (i.e. respiratory CO_2 released per unit biomass produced by growth and by the maintenance of the existing biomass: both, per unit time and unit mass; Penning de Vries, 1975), and W_{live} is the living biomass (Amthor, 2000). W_{live} can be broken down further:

$$W_{\text{live}} = W_{\text{live_woody}} + W_{\text{green}}, \quad (3)$$

where $W_{\text{live_woody}}$ and W_{green} are the biomass of live woody pools (living cells in stem, branches and coarse roots) and nonwoody tissues (leaves and fine roots) respectively. Because plant tissues require N as a component of the enzymes that sustain metabolic processes (including respiration), living biomass is often expressed in nitrogen units, g N/ground area (Cannell & Thornley, 2000), while respiration is expressed in carbon units. Then m_R is in units of $\text{g C g N}^{-1} \text{time}^{-1}$ (Penning de Vries, 1975). Temporal changes in $W_{\text{live_woody}}$ can be summarized by first-order biochemical kinetics:

$$\underbrace{\frac{dW_{\text{live_woody}}}{dt}}_{\text{relative change}} = \underbrace{\varphi \cdot G_{\text{live_woody}}}_{\text{incoming flux}} - \underbrace{W_{\text{live_woody}} \cdot \tau}_{\text{outgoing flux}}, \quad (4)$$


where $G_{\text{live_woody}}$ is the part of G allocated to live woody, φ converts carbon to nitrogen content (g N/g C) and τ is the live woody turnover rate per unit time (t). A similar expression can be written

for W_{green} . The first term on the right-hand side of Equation (4) represents the 'incoming' flux of new living cells; while the second term represents the 'outgoing' flux of living cells that die and become metabolically inactive. But while W_{green} may be only a small fraction of total forest biomass, not changing much after canopy closure, $W_{\text{live_woody}}$ (as also total W) becomes large during forest development and is potentially a strong driver of R (Reich et al., 2008). However interpreted and wherever applied, this general approach including a turnover rate parameter (τ) is equally valid for any mass-, area- or volume-based analyses (Thornley & Cannell, 2000).

Setting $\tau = 1 \text{ year}^{-1}$ in Equation (4) would imply a tight coupling between the previous year's growth and the current year's respiration flux—as suggested by Gifford (2003)—and yields a close approximation to the W98 assumption of a fixed ratio between P_n and P , thus cancelling, on an annual scale, any effect of biomass accumulation. The implication of a 1 year lag between carbon fixation and respiration in woody compounds is consistent with the findings of Amthor (2000), Kagawa, Sugimoto and Maximov (2006a, 2006b), Gough et al. (2008, 2009) Richardson et al. (2015) and Richardson et al. (2013) of a physiological asynchrony by about 1 year between P and growth (and thus on growth and maintenance respiration).

Alternatively, setting $\tau = 0.1 \text{ year}^{-1}$ would imply that most new sapwood cells live for many years, and would closely approximate the R06 assumption of proportionality between R and biomass. Thus, the amount of respiring biomass is regulated by the amount of substrate that is produced each year, forming new sapwood, versus the amount that is converted into nonliving tissues and no longer involved in metabolism; the balance of these processes being controlled by τ (see proofs-of-concept in Figure 1a,b and Table 1, for elaboration).

FIGURE 1 Proofs-of-concept for total R_M (a) and live woody biomass accumulation (b) over the course of forest development (time) and increases in size, assuming different live wood turnover (τ , year^{-1}) rate values, from 1 (W98) to 0.1 year^{-1} (R06) including two intermediate values at 0.5 and 0.7 year^{-1} (e.g. White et al., 2000). $R_{M\text{-green}}$ (i.e. leaf and fine root R_M) was assumed constant over time and arbitrarily equal to 2. Summing up $R_{M\text{-wood}}$ and $R_{M\text{-green}}$ gives the total R_M . Initial woody biomass was arbitrarily considered equal to 10, new annual live wood was also arbitrarily considered equal to 10, $m_R = 0.2$ ($R_M = (W_{\text{live_woody}} + W_{\text{green}}) \times m_R$; see Equation 3). (b) Initial woody biomass was arbitrarily considered equal to 10, new annual live wood was arbitrarily considered equal to 10. The model is: $W_{\text{live_woody}}(t+1) = W_{\text{live_woody}}(t) + \Delta W_{\text{live_woody}}(t+1) - \Delta W_{\text{out_wood}}(t+1)$ (see Equation 4 in the main text) [Colour figure can be viewed at wileyonlinelibrary.com]



τ level	Corresponding underlying assumption	Reference
$\tau = 0.1$	Low turnover rate, which implies only accumulation of respiring biomass (i.e. $R \propto \text{biomass}$)	Reich et al. (2006)
$\tau = 1$	High turnover rate, with death of cells annually equalling live cell production (i.e. $R \propto P$)	Waring et al. (1998)
$0.1 < \tau < 1$	Intermediate turnover rate	e.g. White et al. (2000); see Box 2
$\tau = 0$	Functionally impossible, because it would imply no mortality of cells	—
$\tau > 1$	Physically impossible, because turnover would exceed the number of available living cells	—

TABLE 1 Underlying modelling assumptions adopted in the analysis

BOX 1 The function and dynamics of NSC

Nonstructural carbohydrate is a surprisingly poorly known component of the whole-tree carbon balance, and commonly disregarded in models (Merganičová et al., 2019; Schiestl-Aalto et al., 2019). However, the ability of trees to prioritize storage over growth depends on the role of NSC in allowing temporal asynchrony between carbon demand and carbon supply (Fatichi, Leunzinger, & Körner, 2014). Such imbalances are assumed to be buffered by drawing down NSC reserves. Recent studies support this assumption, showing that during periods of negative carbon balance (for example during the dormant season, periods of stress or natural or artificially induced defoliation episodes) NSC is remobilized and transported from the sites of phloem loading, while during periods of positive carbon balance plants preferentially allocate recently assimilated carbon to replenish NSC. Only afterwards is 'new' carbon used to sustain growth (Huang et al., 2019; Weber et al., 2018). Because ultimately plant survival depends more on metabolic carbon demands than on growth, some have argued that all positive carbon flows should be used to replenish NSC at the expense of growth until a minimum NSC pool size (30%–60% of the seasonal maximum, Martínez-Vilalta et al., 2016) is reached ('active' storage: Sala, Woodruff, & Meinzer, 2012), thus maintaining a safety margin against the risk of carbon starvation (Huang et al., 2019; Wiley & Helliker, 2012). Note that this assumption departs from the notion that NSC is a mere reservoir for excess supply of carbon relative to growth demand ('passive' storage: Kozłowski, 1992). In the model, carbon allocation to all tree structural and nonstructural pools is computed here daily and is controlled by functional constraints due to direct and lagged C-requirements (Huang et al., 2019; Merganičová et al., 2019). It is assumed that a minimum NSC threshold level concentration (11% of sapwood dry mass for deciduous and 5% for evergreen species: Genet, Bréda, & Dufrêne, 2010) has to be maintained for multiple functions including osmoregulation, cell turgor, vascular integrity, tree survival (reviewed in Hartmann & Trumbore, 2016) and organ-specific phenology (leaf and fine-root formation). The greater the sapwood mass, the greater the minimum NSC threshold must be (Dietze et al., 2014). For deciduous trees, four phenological phases are distinguished: (a) the dormant phase, where R is fuelled by NSC-consumption; (b) the leaf onset phase, when leaf and fine root production consume NSC (unless the carbon balance is positive, in which case new assimilates are used) until the predicted maximum annual leaf area index (LAI) is reached; (c) the full growing phase, when new assimilates are allocated to stem, coarse roots, branch and fruits, and only into the NSC pool if this is below its minimum level; (d) the leaf fall phase, when all assimilates are allocated to the NSC reserve pool while some (~10%) NSC is relocated from falling leaves and dying fine roots (Capioli et al., 2013; Collalti et al., 2016). For evergreen species the model follows a simpler schedule consisting of a first maximum growth phase, when the model allocates NSC to foliage and fine roots up to peak LAI, and a second full growing phase, when the model allocates to all of the pools (Kuptz et al., 2011). Such patterns of whole-tree seasonal NSC dynamics have been all recently confirmed by Furze et al. (2019) and Fierravanti, Rossi, Kneeshaw, De Grandpré, and Deslauriers (2019) and a similar phenological and carbon allocation scheme has been adopted by other models (e.g. Arora & Boer, 2005; Krinner et al., 2005).

Because carbon supply (photosynthesis) and carbon metabolic demand (respiration) are not necessarily synchronized, the model assumes that temporary carbon imbalances between P and R (implying $P_n < 0$; Roxburgh, Berry, Buckley, Barnes, & Roderick, 2005) are

met by the remobilization or recycling of NSC stored during previous year(s)—so long as the NSC pool is not completely emptied (the carbon starvation hypothesis; McDowell et al., 2008). A full description of the modelled NSC dynamics is provided in Box 1.

2.2 | Simulation set-up

The logic described above was implemented in a process-based forest growth model (3D-CMCC-BGC), parameterized at site level, and applied, as a case study, to an intensively monitored temperate deciduous forest. Additional model description can be found in Supporting Information, Collalti et al. (2014, 2016, 2018, and references therein), and Marconi, Chiti, Nolè, Valentini, and Collalti (2017).

Very limited data are available on the turnover rate τ of live cells in sapwood, which is often either guessed or inferred by model calibration (e.g. White, Thornton, Running, & Nemani, 2000). We carried out 10 simulations with τ varied in arbitrary 0.1 year⁻¹ steps, from $\tau = 1$ year⁻¹ (100% of turnover, all the previous year live cells of sapwood becomes nonrespiring heartwood in the current year) with R mostly depending on the left-hand side term of Equation (2) ($R \sim g_R G$), down to $\tau = 0.1$ year⁻¹ (only 10% of the previous year's live cells of sapwood biomass dies) and R mostly depending on the right-hand side term of Equation (2) ($R \sim m_R W_{\text{live}}$). Thus, we started with the largest prior distribution for τ , assuming that values outside this range are not functionally possible (Table 1). This approach ensures that any difference in model results reflects difference in specific model assumptions (respiration controlled by photosynthesis or biomass) rather than model structure. We are unaware of any studies reporting changes in τ with age or biomass; we have therefore necessarily assumed that τ is constant in time.

The standard model configuration assigns $\tau = 0.7$ year⁻¹ (Collalti et al., 2019) and this same value has been used by several authors in various modelling contexts (e.g. Bond-Lamberty, Gower, Ahl, & Thornton, 2005; Tatarinov & Cenciala, 2006). Other models have applied different values (see Box 2). Zaehle, Sitch, Smith, and Hatterman (2005), Poulter et al. (2010), and Pappas, Fatichi, Leuzinger, Wolf, and Burlando (2013) found that τ is a critical parameter for both LPJ-DGVM and LPJ-GUESS. We are not aware of similar sensitivity analyses for other models. Leaf and fine root turnover rates are assumed here to be 1 year⁻¹, appropriately for deciduous trees (Pietsch, Hasenauer, & Thornton, 2005). The model parameters accounting for 'age effects' (e.g. those controlling, among other things, leaf conductance: Kirschbaum, 2000; Smith, Prentice, & Sykes, 2001) were set arbitrarily large, to avoid building in prior assumptions. Age- and size-effects are therefore considered synonymous (Mencuccini et al., 2005). A stochastic background whole-tree mortality rate (1% of trees removed each year) was retained and included in Equation (4) to ensure realistic self-thinning (Kirschbaum, 2005; Smith et al., 2001). All other parameters were left unchanged from the standard model configuration.

2.3 | Test site and model run

The model was applied to simulate 150 years of even-aged stand development in a stand of European beech (*Fagus sylvatica* L.; Sorø, Denmark; Wu et al., 2013; Reyer et al., 2019) in daily time steps from 1950 to 2100. The reasons for choosing this stand are: (a) the

BOX 2 Turnover rates and other uncertainties in models

Most vegetation models assume, among other parameters commonly maintained constant, a fixed rate of sapwood turnover, τ . However, lack of information on this parameter has been already shown to be an important source of uncertainty in the modelled carbon balance of vegetation stands (Collalti et al., 2019; Goulden et al., 2011; Malhi, 2012). Values adopted in current models include: $\tau = 0.7$ year⁻¹ in CLM (Oleson et al., 2013), Forest v.5.1 (Schwalm & Ek, 2004), 3D-CMCC-CNR (Collalti et al., 2019) and Biome-BGC (Thornton et al., 2002); $\tau \sim 0.75$ year⁻¹ in CASTANEA (Dufrêne et al., 2005); $\tau = 0.85$ year⁻¹ in LPJ-GUESS (Smith et al., 2001); $\tau = 0.95$ year⁻¹ in SEIB-DGVM (Sato, Itoh, & Kohyama, 2007), LPJ-DGVM (Sitch et al., 2003) and NCAR-LSM (Bonan, Levis, Sitch, Vertenstein, & Oleson, 2003); and $\tau \sim 1$ year⁻¹ in CARAIB (Warnant, François, Strivay, & Gérard, 1994), PnET (Wythers, Reich, & Bradford, 2013) and ORCHIDEE (Krinner et al., 2005).

Additional sources of uncertainty include the lack of consideration of a size- or age-related decline in the ratio of living to dead cells (suggesting a declining τ) (Ceschia, Damesin, Lebaube, Pontailier, & Dufrêne, 2002; Damesin, Ceschia, Le Goff, Ottorini, & Dufrêne, 2002), the effect of changes in climate (which could temporarily increase τ to reduce maintenance costs in favour of growth: Doughty et al., 2015), changes in tissue N and NSC concentrations (Machado & Reich, 2006; Thurner et al., 2017) and, a probable, genetically controlled down-regulation of basal respiration rates with the ageing of cells (Carey, Sala, Keane, & Callaway, 2001; Wiley, Hoch, & Landhäusser, 2017). Moreover, both τ and basal respiration rates (g_R and m_R) are likely to vary among different tree biomass pools (Reich et al., 2008). Respiratory carbon losses per unit plant mass may also change to sustain growth as an acclimatory response to carbon demand due to increasing plant size, and perhaps with changing climate (Smith & Stitt, 2007). These hypotheses are all grounded in theory, but are supported by very limited observations (Friend et al., 2014; Thurner et al., 2017).

extensive literature on European beech, allowing key parameter values to be assigned with confidence; (b) the exceptional quantity and length of data available at the Sorø site for initializing in 1950 and evaluating the model more than 50 years later, thus, allowing long-term processes (including woody biomass accumulation) to emerge; and (c) because the trees are deciduous, we can assume a complete annual turnover of leaves and fine roots and, therefore, more easily disentangle the contributions of W_{green} and $W_{\text{live,woody}}$. Deciduous species are also expected to have greater within-season variability in P_n ; P , and greater asynchrony in carbon supply and demand, than evergreen species (Dietze et al., 2014; Martínez-Vilalta et al., 2016). However, both the model assumptions and its results are based on

general principles and expected to apply more generally than solely to this specific model and site.

We simulated forest development up to 2100, consistent with the common economic rotation length for this species in northern Europe. After canopy closure, modelled LAI and the relative amounts of leaf and fine-root biomass became stable or even decreased slightly, as is usually observed (Yang, He, Aubrey, Zhuang, & Teskey, 2016; Yang, Luo, & Finzi, 2011). Therefore, changes in modeled R , and its components R_M and R_G , could be attributed to changes in the total amount of living woody biomass and the costs of its maintenance.

In 1950, the stand was aged 30 years with an average tree diameter at breast height of ~6 cm and a density of 1,326 trees/ha. Model state variables were initialized using species-specific functional and allometric relationships from the literature, and previous model applications at this site (Collalti et al., 2016, 2018; Marconi et al., 2017). Model sensitivity to parameter values and their uncertainties have been assessed in depth in a previous work (see Collalti et al., 2019, especially their figure 2 and table 3). Management, in the form of thinning, occurred at the site only up to 2014. After that year, only stochastic mortality was accounted for in the model. Live wood was initialized at 15% of sapwood biomass (as the fraction of current year sapwood: Pietsch et al., 2005) and assigned a C:N ratio of 48 g C/g N, not changing with increasing biomass (Ceschia et al., 2002; Damesin, 2003). The minimum concentration of NSC was assumed to be ~11% of sapwood dry mass (Genet et al., 2010; Hoch, Richter, & Körner, 2003; Martínez-Vilalta et al., 2016) consistent with measurements

on deciduous species (and specifically beech). Daily meteorological forcing variables were obtained as historical ensemble means from five Earth System Models up to 2005 provided by the Inter-Sectoral Impact Model Intercomparison Project (ISI-MIP; Warszawski et al., 2014). Data for the period 1995–2005 were then randomly repeated up to 2100. Additional simplifying assumptions were made in order to focus specifically on the effects of increases in tree size, as follows: no disturbances (whether herbivory or management) after 2014; no effect of changes in soil N availability, thus excluding confounding effects of altered N deposition; and, importantly, to avoid possible confounding of temperature effects on R_M with other warming effects, a stable (1995–2005) climate and atmospheric CO_2 concentration (~380 $\mu\text{mol/mol}$). Exports of carbon to exudates and BVOCs are very slight in this species, and they could therefore be neglected.

3 | RESULTS

3.1 | Data-model agreement

The standard model configuration satisfactorily reproduced P , R , P_n and the ratio $P_n:P$ when compared to independent, site-level, carbon balance data (Wu et al., 2013) for the period 2006–2010 (Figure 2; Table 2), corresponding to a stand age of ~85–90 years. P was in agreement with eddy covariance data, while R was slightly underestimated compared to values in Wu et al. Consequently, the model overestimated the average $P_n:P$ ratio by 14% compared to Wu et al. However,





FIGURE 2 Model results for (a) $P_n:P$ ratio (dimensionless), (b) net primary production (P_n , $\text{g C m}^{-2} \text{ year}^{-1}$) and (c) autotrophic respiration (R , $\text{g C m}^{-2} \text{ year}^{-1}$) performed with varying τ (coloured lines). The beginning of simulations correspond to 1950 (stand age 30 years); the end of simulations correspond to 2100 (stand age 180 years). The dark-pointed red line can be considered as a mechanistic representation of W98's fixed $P_n:P$ ratio ($\tau = 1 \text{ year}^{-1}$), while the dark pink line approximates R06's scaling relationship between R and biomass ($\tau = 0.1 \text{ year}^{-1}$). Orange dotted lines represent Amthor's (2000) (A00) 'allowable' range for the $P_n:P$ ratio (0.65–0.2). The red dots give the average measured values (Wu et al., 2013) at the site for (a) $P_n:P$ ratio, (b) P_n and (c) R . Vertical bars represent the standard deviation with horizontal bars representing the period 2006–2010 (stand age ~85–90 years). The shaded area represents the overall uncertainty of model results [Colour figure can be viewed at wileyonlinelibrary.com]

TABLE 2 Model validation (averages for the years 2006–2010), in brackets standard deviation (when available). Literature data come from Wu et al. (2013)

2006–2010 mean (\pm SD)	$P_n:P$	P	R	P_n	$P_{n,woody}$	$P_{n,litter}$	Above- ground-R	Below- ground-R	Above- ground woody stocks	Below- ground woody stocks	Total woody stocks
Units	Unitless	$\text{g C m}^{-2} \text{ year}^{-1}$							g C/m^2		
Literature	0.37	1,881 (± 127)	1,173 (± 143)	708 (± 65)	307 (± 57)	401	872	301	9,885 (± 279)	1,848 (± 160)	11,733 (± 281)
Modelled	0.45 (± 0.02)	1,706 (± 52)	937 (± 30)	768 (± 60)	309 (± 56)	314 (± 9)	635 (± 20)	302 (± 9)	8,993 (± 278)	1,954 (± 545)	10,948 (± 333)

FIGURE 3 Regression analyses between whole-plant autotrophic respiration (R , $\text{g C m}^{-2} \text{ year}^{-1}$) and (a) whole-plant carbon (W ; g C/m^2), (b) carbon in living pools ($W_{\text{live,woody}} + W_{\text{green}}$; g C/m^2), (c) whole-plant nitrogen (W ; g N/m^2) and (d) nitrogen in living pools ($W_{\text{live,woody}} + W_{\text{green}}$; g N/m^2). Different colours represent different τ values as described in the legend panels (with $\tau = 0.1 \text{ year}^{-1}$, $n = 117$; with $\tau = 0.2 \text{ year}^{-1}$, $n = 149$; otherwise $n = 150$) [Colour figure can be viewed at wileyonlinelibrary.com]

Wu et al. argued that the values of R they obtained (by subtracting modelled heterotrophic from measured ecosystem respiration) may have been overestimated, given also the large standard deviation ($\pm 143 \text{ g C m}^{-2} \text{ year}^{-1}$). The model results are otherwise in good agreement with Wu et al. for woody carbon stocks (both above- and below-ground), annual wood production (the sum of carbon allocated to stems, branches and coarse roots), and annual above- and below-ground litter production (the sum of carbon allocated to leaves and fine roots; Table 2). Modelled respiration of the woody compartments, leaf and total (above- and below-ground) respiration, and NSC pool and fluxes, are all compatible with values reported by previous investigations, and within the ranges of total, wood and leaf respiration, and $P_n:P$ ratios reported for European beech (e.g. Barbaroux & Br  da, 2002; Barbaroux, Br  da, & Dufr  ne, 2002; Davi, Barbaroux, Francois, & Dufr  ne, 2009; Genet et al., 2010; Granier, Breda, Longdoz, Gross, &

Ngao, 2008; Guidolotti, Rey, D'Andrea, Matteucci, & De Angelis, 2013; Knohl, Schulze, Kolle, & Buchmann, 2003). A model validation forced by actual measured climate at this site is also described in previous papers (Collalti et al., 2016, 2018; Marconi et al., 2017).

3.2 | The effect of varying τ

The simulations produced a spectrum of diverging trajectories, ranging from an approximately steady-state with constant $P_n:P$ ratio (for large τ) to a constantly decreasing $P_n:P$ ratio (for small τ ; Figure 1a). For $\tau = 1 \text{ year}^{-1}$, $P_n:P$ stays close to 0.5. For $\tau \leq 0.2 \text{ year}^{-1}$, $P_n:P$ eventually falls below the lower limit of commonly observed values (0.22; Collalti & Prentice, 2019) and the physiological limit of 0.2 proposed by Amthor (2000). Figure 2 also shows the effects of varying τ in determining different trajectories for P_n (Figure 2b)

and R (Figure 2c) and consequent differences in the partitioning between R_M and R_G (Figure S2). Modelled R , at the end of simulations, ranging from $\sim 800 \text{ g C m}^{-2} \text{ year}^{-1}$, giving $P_n \sim 900 \text{ g C m}^{-2} \text{ year}^{-1}$ and $P \sim 1,700 \text{ g C m}^{-2} \text{ year}^{-1}$ (Figure S1 and for NSC flux Figure S3b) consistent with a steady-state between R or P_n and P , to two cases ($\tau = 0.1, 0.2 \text{ year}^{-1}$) in which trees die from starvation.

The model did not generate any consistent power-law relationship between R and biomass either for $b \sim 1$ (i.e. R_0), or for $\sim 3/4$ (Mori et al., 2010), or for $\sim 2/3$ (Makarieva, Gorshkov, & Li, 2005) or for $\sim 3/5$ (Michalet et al., 2014; Table S1). The simulations indicated $b \sim 1$ initially, shifting with increasing tree size to $b \sim 0.74$ for $\tau = 0.1 \text{ year}^{-1}$ ($R^2 = .99$; $n = 117$) or 0.19 for $\tau = 1 \text{ year}^{-1}$ ($R^2 = .84$; $n = 150$; 'n' corresponds to years of simulation). For the relation between R and whole-plant N, again the simulations indicated $b \sim 1$ initially, shifting to $b \sim 0.82$ for $\tau = 0.1 \text{ year}^{-1}$ ($R^2 = .99$; $n = 117$) or 0.27 for $\tau = 1 \text{ year}^{-1}$ ($R^2 = .82$; $n = 150$; Figure 3a,c). The highest b values corresponded to simulations which ended because the trees died.

4 | DISCUSSION

4.1 | R is not entirely determined by P

A constant $P_n:P$ ratio, as implied by W98's hypothesis and obtained here by setting $\tau = 1 \text{ year}^{-1}$, conflicts with observations from many different tree species that show a substantially slower turnover rate of living cells. In fact, parenchyma cells within secondary xylem are very often more than a year old, and can be up to 200 years old (Spicer & Holbrook, 2007). The constant ratio hypothesis is also contrary to the evidence in trees that much of the recently fixed assimilate pool is at first stored as reserves, and only later used for metabolism or growth (Schiestl-Aalto, Kulmala, Nikinmaa, & Mäkelä, 2015; Schiestl-Aalto et al., 2019). Indeed, there are some reports of decoupling between growth (which would imply some CO_2 released for both R_G and subsequently R_M) and photosynthesis—with growth ceasing long before photosynthesis—because of the different sensitivities of growth and photosynthesis to environmental drivers. Kagawa, Sugimoto, and Maximov (2006a) reported for *Larix gmelinii* Mayr. that up to 43%, and according to Gough, Flower, Vogel, Dragoni, and Curtis (2009) up to 66%, of annual photosynthates in bigtooth aspen (*Populus grandidentata* Michx.) and northern red oak (*Quercus rubra* L.) are used during the year(s) after they have been fixed. Gaudinski et al. (2009), Malhi (2012) and Delpierre, Berveiller, Granda, and Dufrêne (2016) all found negative correlations between annual carbon inflows and above- or below-ground wood growth, from temperate to tropical tree species. Analysing Luyssaert et al.'s (2007) global database, Chen, Chen, Yang, and Robinson (2013) found that R does not scale isometrically with P . Some authors have suggested that R_G could be supplied exclusively by recent photosynthates while R_M by previously stored ones (Lotscher, Klumpp, & Schnyder, 2004). Along the same lines, Maier, Johnsen, Clinton and Ludovici (2010) for loblolly pine trees (*Pinus taeda* L.), Kuptz et al. (2011) for beech (*F. sylvatica*

L.) and Norway spruce (*Picea abies* Karst.) and Lynch, Matamala, Iversen, Norby, and Gonzalez-Meler (2013) for a sweetgum plantation (*Liquidambar styraciflua* L.) found that both R_G and R_M are not completely satisfied by recent assimilates, and that some current R_M can be derived from woody tissues constructed in previous years. Litton, Raich, and Ryan (2007) and Yang et al. (2016) both found low correlations between respiration and the annual production of woody compounds in large data sets. Many studies have also reported little variation in the CO_2 efflux from sapwood in relation to tree-ring age, despite a stepwise decrease in the fraction of living cells towards the centre of the stems (e.g. Ceschia et al., 2002; Pallardy, 2010; Spicer & Holbrook, 2007). These various observations imply that some carbon is fixed 1 year and used for the tree's own growth and metabolism in the next or subsequent years, and that the inner sapwood contains a population of living cells formed in previous years.

These observations are all incompatible with the hypothesis of a tight coupling of R and P (alone), and with model results obtained by assuming complete turnover of live cells in sapwood during a single year.

4.2 | R is not entirely determined by biomass

On the other side of the ledger, model simulations indicate that low τ values ($\leq 0.2 \text{ year}^{-1}$) can lead to excessively high respiration burdens, impossibly low $P_n:P$ ratios (< 0.2) and ultimately carbon starvation when all NSC is consumed and whole-tree R_M or growth can no longer be sustained (Figure 2). This model result is quantitatively dependent on the values adopted for C:N ratio and the minimum NSC-pool which increases with tree size, but it is consistent with the idea that $P_n:P$ ratios ≤ 0.2 are not physiologically sustainable (Amthor, 2000). Amthor described, for a large data set comprising grasses, tree crops and forest trees worldwide, the 0.65–0.2 bounds as reflecting maximum growth with minimum maintenance expenditure (0.65) and minimum growth with maximum physiologically sustainable maintenance costs (0.2). Such a minimum $P_n:P$ value agrees also with Keith et al.'s (2010) reasoning (analysing *Eucalyptus* forests of south-eastern Australia) that trees always require some annual biomass production in order to survive. With such low τ , simulated woody $R_M:R$ exceeds 90%, a value much higher than those (56%–65%) reported by Amthor and Baldocchi (2001; Figure S2). This situation initiates a spiral of decline, whereby neither P nor NSC drawdown are sufficient to avoid a long-term carbon imbalance (Figure S3a,b; Weber et al., 2018; Wiley et al., 2017).

Slightly higher τ values (from 0.3 to 0.5 year^{-1}) were found to limit woody biomass increase because of high NSC demand, leading to a shift in the allocation of assimilates to refill NSC at the expense of growth, and $P_n:P$ values close to 0.2 (Figure 2a). Values of $\tau > 0.5$ did not show such behaviour and allowed structural and nonstructural compounds to accumulate in parallel, while P_n gradually declined and eventually levelled-off. This scenario allows structural biomass accumulation to continue even in older trees, as has been observed (Stephenson et al., 2014).

4.3 | Scaling relationships

We simulated forest dynamics from juvenile up to very large, mature trees while R06's results supporting isometric scaling were based on measurements of seedlings and 6- to 25-year-old trees with, presumably, very little heartwood. Some other studies have suggested that the scaling slope of approximately 1 for whole-plant mass may be valid early in stand development, but that the exponent may eventually become smaller than 3/4, a phenomenon that has been termed 'ontogenic drift' (Makarieva et al., 2008). Piao et al. (2010) in a global analysis also found a low correlation, and a low scaling exponent, between R and whole-plant biomass ($b = 0.21$, corresponding to $\tau \sim 0.9$ in our simulations). Piao et al. (2010) argued that, for large mature trees, an increasing fraction of woody C and N biomass is composed of metabolically inactive heartwood, and concluded that a linear relationship between respiration and whole-plant biomass should not be expected (even if there is a linear relationship of respiration to the live component of woody biomass), while a curvilinear relationship at the small end of the size-spectrum seemed more appropriate (Kozłowski & Konarzewski, 2005). Li, Han, and Wu (2005) also found no evidence for an isometric or 3/4 power scaling relationship, indicating instead a range between 1.14 and 0.40, decreasing with plant size. The only approximately isometric relationship we found in our simulations—across all τ used—was between R and the living components of biomass C (b in the range of 0.8–1, with R^2 always $>.93$) and biomass N ($b \sim 0.9$, with R^2 always $>.97$; Figure 3b,d; Gruber, Wieser, & Oberhuber, 2009; Kerkhoff & Enquist, 2005; Makarieva et al., 2005). Conversely, and in accordance with Piao et al. (2010), by considering all woody biomass (sapwood and heartwood), b consistently deviates from linearity for both C and N in biomass, because—as observed in mature and big trees—an increasing amount of biomass is composed of metabolically inactive tissues that do not respire.

None of these findings are compatible with a tight isometric relationship of R to whole-plant C (or N) biomass as proposed by R06.

4.4 | R is determined by P , biomass and the demand for reserves

Plants store large amounts of NSC (potentially enough to rebuild the whole leaf canopy one to more than four times: Hoch et al., 2003) and, when needed, plants can actively buffer the asynchronies between carbon demand (i.e. R and G) and supply (i.e. P) by tapping the pool of nonstructural carbon (see Figure S3 for NSC trends). Several lines of evidence and a growing body of literature support the view of an active sink of NSC. That is, NSC competes with growth, while it controls R (and including other nonmetabolic functions, see Hartmann & Trumbore, 2016), in a compensatory mechanism (high NSC demands for respiration means low carbon supply for biomass growth and vice versa). Schuur and Trumbore (2006) and Carbone, Czimizik, McDuffee, and Trumbore (2007) for boreal black spruce forest (*Picea mariana* B. S. P), and Lynch et al. (2013) for a *L. styraciflua* plantation, all reported that plant-respired CO_2 is a mixture of old and new assimilated carbohydrates. Likewise, Vargas, Trumbore, and Allen (2009) for semideciduous tree species, Carbone et al. (2013), Richardson et al. (2015) and Richardson et al.

(2013) for red maple trees (*Acer rubrum* L.), Muhr et al. (2013, 2016) for different Amazonian tree species and Solly et al. (2018) for pines (*Pinus sylvestris* L.), beeches (*F. sylvatica* L.), spruces (*P. abies* Karst) and birches (*Betula nana* L.), all found that old NSC (up to 17 years old) remobilized from parenchyma cells can be used for growth or metabolism.

Aubrey and Teskey (2018) found that carbon-starved roots and whole-tree saplings die before complete NSC depletion in longleaf pine (*Pinus palustris* L.), but the threshold NSC level at which this happens remains unknown for most species. These thresholds are likely to vary among tissues (Weber et al., 2018), species (Hoch et al., 2003), phenotypes, habit and wood anatomy (Dietze et al., 2014), and increase with tree size (Sala et al., 2012). Others have reported that aspen trees (*Populus tremuloides* Michx) cannot draw down NSC to zero because of limitations in carbohydrate remobilization and/or transport (Wiley et al., 2017). A minimum NSC level, which has been found to proportionally increase with biomass, may also be required to maintain a safety margin and a proper internal functioning of trees (including osmoregulation), regardless of whether growth is limited by carbon supply (Huang et al., 2019; Martínez-Vilalta et al., 2016; Sala, Fouts, & Hoch, 2011; Sala et al., 2012; Woodruff & Meinzer, 2011). Genet et al. (2010) found for beech and sessile oak (*Quercus petraea* [Matt.] Liebl.) shifts during ontogeny in carbon allocation from biomass growth to reserves regardless of seasonal fluctuations, habitat and climate. Palacio, Hernández, Maestro-Martínez, and Camarero (2012) found that black pine trees (*Pinus nigra* Arnold) that were repeatedly defoliated for 11 years, and left to recover for another 6 years, showed reduced growth but similar stem NSC concentration when compared to control trees. Fierravanti et al. (2019) found that low NSC accumulation in conifers defoliated by spruce budworm led to a reduction in growth and an increase in mortality.

It has further been suggested that a considerable fraction of NSC (mostly starch) in the inner part of wood may become compartmentalized and sequestered away from sites of phloem loading, and thus is no longer accessible for either tissue growth or respiration (Sala et al., 2012). Root exudation to mycorrhizal fungi and secondary metabolites (not accounted for here) could also accelerate NSC depletion (Pringle, 2016), and potentially create a risk of carbon starvation even for values of τ well above 0.2.

Overall, asynchrony between (photosynthetic) source and (utilization) sink implies some degree of uncoupling of R , and consequently P_n (and growth), from P and biomass. Carbon demand for metabolism and growth can be mediated by tapping the pool of NSC but only to the extent and to the amount that it is accessible and useable by plants. Therefore, if this active role of NSC can be experimentally confirmed, it will imply that plants prioritize carbon allocation to NSC over growth.

4.5 | Implications

It has been suggested that the observed decline of P_n during stand development cannot be exclusively caused by increasing respiration costs with tree size (Tang, Luyssaert, Richardson, Kutsch, & Janssens, 2014). The idea, implicit in the growth and maintenance respiration paradigm—that the maintenance of existing biomass (R_M) is a 'tax' that must be

paid first and which ultimately controls growth—has also been criticized for lack of empirical support (Gifford, 2003). While this paradigm has some weaknesses (Thornley, 2011), and has not changed much over the last 50 years despite some theoretical and experimental refinements (e.g. accounting for temperature acclimation: Tjoelker et al., 1999), it reflects the prevailing assumption embedded in models because, so far, no other general (and similarly promising) mechanistic approach to the modelling of whole-plant respiration has been proposed.

Although plant physiologists are well aware that respiration is actually neither entirely determined by photosynthesis nor entirely determined by biomass, but rather by plants' energy requirements for their functioning and growth, we highlight the persistent large uncertainty surrounding this issue in the forestry and forest ecology literature. Both the literature reviewed here and our model results show that any successful modelling approach for plant respiration must necessarily allow plants to steer a middle course between tight coupling to photosynthesis (inconsistent with a carbon steady-state in forest development, and with many observations) and dependence on ever-increasing biomass (risking carbon starvation and death), coupled to the buffering capacity of reserves during carbon imbalances (see Box 1). It seems likely that plants strive to keep an appropriate quantity of living cells that can effectively be sustained by photosynthesis or, when necessary, by drawing on NSC and down-regulating allocation to nonphotosynthetic, but metabolically active, tissues as to minimize maintenance costs (Makarieva et al., 2008). This would suggest active control on carbon use efficiency and on the turnover of the living cells by plants. Yet, despite its importance, NSC use is overlooked in 'state-of-the-art' vegetation models. The present study has not been able to provide tight numerical constraints on τ . However, we can unequivocally reject the two, mutually incompatible simplifying hypotheses as both conflict with a large and diverse body of evidence.

Other processes, including hydraulic and nutrient limitations, may be in play (Carey et al., 2001; Xu et al., 2012). Malhi et al. (2015) argued for a link between high whole-plant mortality rates and high forest productivity as ecophysiological strategies that favour rapid growth may also result in fast turnover of trees. However, Spicer and Holbrook (2007) noted that metabolic activity does not decline with cell age; and Mencuccini et al. (2005) noted that effects of age per se (including cellular senescence and apoptosis) are likely not responsible for declining P , but are linked to the functional and structural consequences of increasing plant size. This is an important conclusion because it allows models to avoid accounting explicitly for age.

In conclusion, to reduce the large uncertainty surrounding this issue, it will be necessary on the one hand to use models that explicitly account for the turnover of biomass and the reserves usage; and on the other hand, to carry out experimental and field measurements of the dynamics of living cells in wood and the availability of and demand for labile carbon stores. These processes have a direct bearing on the stocks and fluxes that drive the carbon balance of forests.

ACKNOWLEDGEMENTS

We gratefully acknowledge Owen K. Atkin, Dennis Baldocchi, Ben Bond-Lamberty, Marcos Fernández-Martínez, Roger Gifford, Francesco Loreto, Maurizio Mencuccini, Joe Landsberg and Richard Waring, including the two reviewers, for reading the many drafts of this paper and having returned many insightful and constructive comments. We thank Andreas Ibrom for providing the site data and for the useful discussions and Carlo Trotta, Angelo Rita, Giulia Mengoli, Elisa Grieco for technical assistance. This work contributes to the AXA Chair Programme in Biosphere and Climate Impacts and the Imperial College initiative on Grand Challenges in Ecosystems and the Environment. It has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No: 787203 REALM). We thank the ISI-MIP project (<https://www.isimip.org/>) and the COST-Action PROFOUND (FP 1304) for providing the climate historical scenarios and site data used in this work. The 3D-CMCC-FEM and BGC model code are freely available at: <https://github.com/3D-CMCC-CNR-FEM>

AUTHOR CONTRIBUTIONS

A.C., M.T., G.H., G.G., G.P., and I.C.P. designed the study. A.C., G.H., G.G., and M.G.R. carried out modelling work and data analysis. A.C., G.G., M.G.R., G.P., and I.C.P. drafted the manuscript with contributions from all authors.

ORCID

Alessio Collalti  <https://orcid.org/0000-0002-4980-8487>

Mark G. Tjoelker  <https://orcid.org/0000-0003-4607-5238>

Günter Hoch  <https://orcid.org/0000-0003-0985-9746>

Annikki Mäkelä  <https://orcid.org/0000-0001-9633-7350>

Gabriele Guidolotti  <https://orcid.org/0000-0002-2052-6523>

Mary Heskel  <https://orcid.org/0000-0003-3227-2978>

Giai Petit  <https://orcid.org/0000-0002-6546-7141>

Michael G. Ryan  <https://orcid.org/0000-0002-2500-6738>

Giovanna Battipaglia  <https://orcid.org/0000-0003-1741-3509>

Giorgio Matteucci  <https://orcid.org/0000-0002-4790-9540>

Iain Colin Prentice  <https://orcid.org/0000-0002-1296-6764>

REFERENCES

- Agutter, P. S., & Tuszyński, J. A. (2011). Analytic theories of allometric scaling. *The Journal of Experimental Biology*, 214, 1055–1062. <https://doi.org/10.1242/jeb.054502>
- Amthor, J. (2000). The McCree-de Wit-Penning de Vries-Thornley respiration paradigms: 30 Years later. *Annals of Botany*, 86, 1–20. <https://doi.org/10.1006/anbo.2000.1175>

- Amthor, J., & Baldocchi, D. (2001). Terrestrial higher plant respiration and net primary production. In J. Roy, B. Sauger, & H. A. Mooney (Eds.), *Terrestrial global productivity* (pp. 33–59). San Diego, CA: Academic Press.
- Arora, V., & Boer, G. (2005). A parameterization for leaf phenology in the terrestrial ecosystem component of climate models. *Global Change Biology*, 11(1), 39–59. <https://doi.org/10.1111/j.1365-2486.2004.00890.x>
- Aubrey, D. P., & Teskey, R. O. (2018). Stored carbohydrates can maintain root respiration for extended periods. *New Phytologist*, 218, 142–152. <https://doi.org/10.1111/nph.14972>
- Barbaroux, C., Brèda, E., & Dufrêne, E. (2002). Distribution of above-ground and below-ground carbohydrate reserves in adult trees of two contrasting broad-leaved (*Quercus cerris* and *Fagus sylvatica*). *New Phytologist*, 157, 605–615. <https://doi.org/10.1046/j.1469-8137.2003.00681.x>
- Barbaroux, C., & Brèda, N. (2002). Contrasting distribution and seasonal dynamics of carbohydrate reserve in stem wood of adult ring-porous sessile oak and diffuse-porous beech trees. *Tree Physiology*, 22, 1201–1210. <https://doi.org/10.1093/treephys/22.17.1201>
- Bonan, G. B., Levis, S., Sitch, S., Vertenstein, M., & Oleson, K. W. (2003). A dynamic global vegetation model for use with climate models: Concepts and description of simulated vegetation dynamics. *Global Change Biology*, 9, 1543–1566. <https://doi.org/10.1046/j.1365-2486.2003.00681.x>
- Bond-Lamberty, B., Gower, S., Ahl, D. E., & Thornton, P. E. (2005). Reimplementation of the Biome-BGC model to simulate successional change. *Tree Physiology*, 25, 413–424. <https://doi.org/10.1093/treephys/25.4.413>
- Campioli, M., Verbeeck, H., Van den Bossche, J., Wu, J., Ibrom, A., D'Andrea, E., ... Granier, A. (2013). Can decision rules carbon allocation for years with contrasting and extreme weather conditions? A case study for three temperate beech forests. *Ecological Modelling*, 263, 42–55. <https://doi.org/10.1016/j.ecolmodel.2013.04.012>
- Cannell, M., & Thornley, J. (2000). Modelling the components of plant respiration: Some guiding principles. *Annals of Botany*, 85, 45–54. <https://doi.org/10.1006/anbo.1999.0996>
- Carbone, M. S., Czimczik, C. I., Keenan, T. F., Murakami, P. F., Pederson, N., Schaberg, P. G., ... Richardson, A. D. (2013). Age, allocation and availability of non-structural carbon in mature red maple trees. *New Phytologist*, 200, 1145–1155. <https://doi.org/10.1111/nph.12448>
- Carbone, M. S., Czimczik, C. I., McDuffee, K. E., & Trumbore, S. E. (2007). Allocation and residence time of photosynthetic products in a boreal forest using a low-level ^{14}C pulse-chase labeling technique. *Global Change Biology*, 13, 466–477. <https://doi.org/10.1111/j.1365-2486.2006.01300.x>
- Carey, E. V., Sala, A., Keane, R., & Callaway, R. M. (2001). Are old forests underestimated as global carbon sinks? *Global Change Biology*, 7, 339–344. <https://doi.org/10.1046/j.1365-2486.2001.00418.x>
- Ceschia, E., Damesin, C., Lebaube, S., Pontailler, J. Y., & Dufrêne, É. (2002). Spatial and seasonal variations in stem respiration of beech trees (*Fagus sylvatica*). *Annals of Forest Science*, 59, 801–812. <https://doi.org/10.1051/forest:2002078>
- Chapin, F. S., Woodwell, G. M., Randerson, J. T., Rastetter, E. B., Lovett, G. M., Baldocchi, D. D., ... Schulze, E.-D. (2006). Reconciling carbon-cycle concepts, terminology, and methods. *Ecosystems*, 9(7), 1041–1050. <https://doi.org/10.1007/s10021-005-0105-7>
- Chen, G., Chen, G., Yang, Y., & Robinson, D. (2013). Allocation of gross primary production in forest ecosystems: Allometric constraints and environmental responses. *New Phytologist*, 200, 1176–1186. <https://doi.org/10.1111/nph.12426>
- Collalti, A., Marconi, S., Ibrom, A., Trotta, C., Anav, A., D'andrea, E., ... Grünwald, T. (2016). Validation of 3D-CMCC Forest Ecosystem Model (v.5.1) against eddy covariance data for 10 European forest sites. *Geoscientific Model Development*, 9, 479–504. <https://doi.org/10.5194/gmd-9-479-2016>
- Collalti, A., Perugini, L., Santini, M., Chiti, T., Nolè, A., Matteucci, G., & Valentini, R. (2014). A process-based model to simulate growth in forests with complex structure: Evaluation and use of 3D-CMCC Forest Ecosystem Model in a deciduous forest in Central Italy. *Ecological Modelling*, 272, 362–378. <https://doi.org/10.1016/j.ecolmodel.2013.09.016>
- Collalti, A., & Prentice, I. C. (2019). Is NPP proportional to GPP? Waring's hypothesis twenty years on. *Tree Physiology*, 39, 1473–1483. <https://doi.org/10.1093/treephys/tpz034>
- Collalti, A., Thornton, P. E., Cescatti, A., Rita, A., Borghetti, M., Nolè, A., ... Matteucci, G. (2019). The sensitivity of the forest carbon budget shifts across processes along with stand development and climate change. *Ecological Applications*, 29(2), 1–18. <https://doi.org/10.1002/eap.1837>
- Collalti, A., Trotta, C., Keenan, T. F., Ibrom, A., Bond-Lamberty, B., Grote, R., ... Anav, A. (2018). Thinning can reduce losses in carbon use efficiency and carbon stocks in managed forests under warmer climate. *Journal of Advances in Modeling Earth System*, 10(10), 2427–2452. <https://doi.org/10.1029/2018ms001275>
- Damesin, C. (2003). Respiration and photosynthesis characteristics of current-year stems of *Fagus sylvatica*: From the seasonal pattern to an annual balance. *New Phytologist*, 158(3), 465–475. <https://doi.org/10.1046/j.1469-8137.2003.00756.x>
- Damesin, C., Ceschia, E., Le Goff, N., Ottorini, J.-M., & Dufrêne, E. (2002). Stem and branch respiration of beech: From tree measurements to estimations at the stand level. *New Phytologist*, 153, 159–172. <https://doi.org/10.1046/j.0028-646X.2001.00296.x>
- Davi, H., Barbaroux, C., Francois, C., & Dufrêne, E. (2009). The fundamental role of reserves and hydraulic constraints in predicting LAI and carbon allocation in forests. *Agricultural and Forest Meteorology*, 149, 349–361. <https://doi.org/10.1016/j.agrformet.2008.08.014>
- Delpierre, N., Berveiller, D., Granda, E., & Dufrêne, E. (2016). Wood phenology, not carbon input, controls the interannual variability of wood growth in a temperate oak forest. *New Phytologist*, 210, 459–470. <https://doi.org/10.1111/nph.13771>
- Dietze, M., Sala, A., Carbone, M. S., Czimczik, C. I., Mantoosh, J. A., Richardson, A. D., & Vargas, R. (2014). Nonstructural carbon in woody plants. *Annual Review of Plant Biology*, 65, 667–687. <https://doi.org/10.1146/annurev-arplant-050213-040054>
- Doughty, C. E., Metcalfe, D., Girardin, C., Amézquita, F. F., Cabrera, D. G., Huasco, W. H., ... Feldpausch, T. R. (2015). Drought impact on forest carbon dynamics and fluxes in Amazonia. *Nature*, 519, 78–82. <https://doi.org/10.1038/nature14213>
- Dufrêne, E., Davi, E., François, C., Le Maire, G., Le Dantec, V., & Granier, A. (2005). Modelling carbon and water cycles in a beech forest. Part I: Model description and uncertainty analysis on modelled NEE. *Ecological Modelling*, 185, 407–436. <https://doi.org/10.1016/j.ecolmodel.2005.01.004>
- Fatichi, S., Leunzinger, S., & Körner, C. (2014). Moving beyond photosynthesis: From carbon source to sink-driven vegetation modeling. *New Phytologist*, 201, 1086–1095. <https://doi.org/10.1111/nph.12614>
- Fierravanti, A., Rossi, S., Kneeshaw, D., De Grandpré, L., & Deslauriers, A. (2019). Low non-structural carbon accumulation in spring reduces growth and increases mortality in conifers defoliated by spruce budworm. *Frontiers in Forests and Global Change*, 2, 15. <https://doi.org/10.3389/ffgc.2019.00015>
- Friedlingstein, P., Meinshausen, M., Arora, V. K., Jones, C. D., Anav, A., Liddicoat, S. K., & Knutti, R. (2014). Uncertainties in CMIP5 climate projections due to carbon cycle feedbacks. *Journal of Climate*, 27, 511–526. <https://doi.org/10.1175/jcli-d-12-00579.1>
- Friend, A. D., Lucht, W., Rademacher, T. T., Keribin, R., Betts, R., Cadule, P., ... Ito, A. (2014). Carbon residence time dominates uncertainty in terrestrial vegetation responses to future climate and atmospheric

- CO₂. *Proceedings of the National Academy of Sciences of the United States of America*, 111(9), 3280–3285. <https://doi.org/10.1073/pnas.1222477110>
- Furze, M. E., Huggett, B. A., Aubrecht, D. M., Stolz, C. D., Carbone, M. S., & Richardson, A. D. (2019). Whole-tree nonstructural carbohydrate storage and seasonal dynamics in five temperate species. *New Phytologist*, 221, 1466–1477. <https://doi.org/10.1111/nph.15462>
- Gaudinski, J. B., Torn, M. S., Riley, W. J., Swanston, C., Trumbore, S. E., Joslin, J. D., ... Hanson, P. J. (2009). Use of stored carbon reserves in growth of temperate tree roots and leaf buds: Analyses using radiocarbon measurements and modelling. *Global Change Biology*, 15, 992–1014. <https://doi.org/10.1111/j.1365-2486.2008.01736.x>
- Genet, H., Bréda, N., & Dufrêne, E. (2010). Age-related variation in carbon allocation at tree and stand scales in beech (*Fagus sylvatica* L.) and sessile oak (*Quercus petraea* (Matt.) Liebl.) using a chronosequence approach. *Tree Physiology*, 30, 177–192. <https://doi.org/10.1093/treephys/tpp105>
- Gifford, R. (2003). Plant respiration in productivity models: Conceptualisation, representation and issues for global terrestrial carbon-cycle research. *Functional Plant Biology*, 30, 171–186. <https://doi.org/10.1071/fp02083>
- Gough, C. M., Flower, C. E., Vogel, C. S., Dragoni, D., & Curtis, P. S. (2009). Whole-ecosystem labile carbon production in a north temperate deciduous forest. *Agricultural and Forest Meteorology*, 149, 1531–1540. <https://doi.org/10.1016/j.agrformet.2009.04.006>
- Gough, C. M., Vogel, C. S., Schmid, H. P., Su, H. B., & Curtis, P. S. (2008). Multi-year convergence of biometric and meteorological estimates of forest carbon storage. *Agricultural and Forest Meteorology*, 148, 158–170. <https://doi.org/10.1016/j.agrformet.2007.08.004>
- Goulden, M. L., McMillan, A. M., Winston, G. C., Rocha, A. V., Manies, K. L., Harden, J. W., & Bond-Lamberty, B. P. (2011). Patterns of NPP, GPP, respiration, and NEP during boreal forest succession. *Global Change Biology*, 17(2), 855–871. <https://doi.org/10.1111/j.1365-2486.2010.02274.x>
- Granier, A., Breda, N., Longdoz, B., Gross, P., & Ngao, J. (2008). Ten years of fluxes and stand growth in a young beech forest at Hesse. *Annals of Forest Science*, 65, 704. <https://doi.org/10.1051/forest:2008052>
- Gruber, A., Wieser, G., & Oberhuber, W. (2009). Intra-annual dynamics of stem CO₂ efflux in relation to cambial activity and xylem development in *Pinus cembra*. *Tree Physiology*, 26, 641–649. <https://doi.org/10.1093/treephys/tpp001>
- Guidolotti, G., Rey, A., D'Andrea, E., Matteucci, G., & De Angelis, P. (2013). Effect of environmental variables and stand structure on ecosystem respiration components in a Mediterranean beech forest. *Tree Physiology*, 33, 960–972. <https://doi.org/10.1093/treephys/tpt065>
- Hartmann, H., & Trumbore, S. (2016). Understanding the roles of non-structural carbohydrates in forest trees – From what we can measure to what we want to know. *New Phytologist*, 211, 386–403. <https://doi.org/10.1111/nph.13955>
- Heskel, M. A., O'Sullivan, O. S., Reich, P. B., Tjoelker, M. G., Weerasinghe, L. K., Penillard, A., ... Atkin, O. K. (2016). Convergence in the temperature response of leaf respiration across biomes and plant functional types. *Proceedings of the National Academy of Sciences of the United States of America*, 113, 3832–3837. <https://doi.org/10.1073/pnas.1520282113>
- Hoch, G., Richter, A., & Körner, C. (2003). Non-structural carbon compounds in temperate forest trees. *Plant, Cell & Environment*, 26, 1067–1081. <https://doi.org/10.1046/j.0016-8025.2003.01032.x>
- Huang, J., Hammerbacher, A., Weinhold, A., Reichelt, M., Gleixner, G., Behrendt, T., ... Hartmann, H. (2019). Eyes on the future – Evidence for trade-offs between growth storage and defense in Norway spruce. *New Phytologist*, 222, 144–158. <https://doi.org/10.1111/nph.15522>
- Huntingford, C., Atkin, O. K., Martinez-de la Torre, A., Mercado, L. M., Heskel, M. A., Harper, A. B., ... Malhi, Y. (2017). Implications of improved representations of plant respiration in a changing climate. *Nature Communications*, 8. <https://doi.org/10.1038/s41467-017-01774-z>
- Kagawa, A., Sugimoto, A., & Maximov, T. C. (2006a). Seasonal course of translocation, storage and remobilization of ¹³C pulse-labeled photoassimilate in natural growing *Larix gmelinii* saplings. *New Phytologist*, 171, 793–804. <https://doi.org/10.1111/j.1469-8137.2006.01780.x>
- Kagawa, A., Sugimoto, A., & Maximov, T. C. (2006b). ¹³CO₂ pulse-labeling of photoassimilates reveals carbon allocation within and between tree rings. *Plant, Cell & Environment*, 29, 1571–1584. <https://doi.org/10.1111/j.1365-3040.2006.01533.x>
- Keith, H., Mackey, B., Berry, S., Lindenmayer, D., & Gibbons, P. (2010). Estimating carbon carrying capacity in natural forest ecosystems across heterogeneous landscapes: Addressing sources of error. *Global Change Biology*, 16, 2971–2989. <https://doi.org/10.1111/j.1365-2486.2009.02146.x>
- Kerkhoff, A. J., & Enquist, B. J. (2005). Ecosystem allometry: The scaling of nutrients stocks and primary productivity across plant communities. *Ecology Letters*, 9, 419–427. <https://doi.org/10.1111/j.1461-0248.2006.00888.x>
- Kira, T., & Shidei, T. (1967). Primary production and turnover of organic matter in different forest ecosystems of the Western Pacific. *Japanese Journal of Ecology*, 17, 70–87. https://doi.org/10.18960/seitai.17.2_70
- Kirschbaum, M. U. F. (2000). Forest growth and species distribution in a changing climate. *Tree Physiology*, 20, 309–322. <https://doi.org/10.1093/treephys/20.5-6.309>
- Kirschbaum, M. U. F. (2005). A model analysis of the interaction between forest age and forest responsiveness to increasing CO₂ concentration. *Tree Physiology*, 25, 953–963. <https://doi.org/10.1093/treephys/25.7.953>
- Knohl, A., Schulze, E.-D., Kolle, O., & Buchmann, N. (2003). Large carbon uptake by an unmanaged 250-year-old deciduous forest in Central Germany. *Agricultural and Forest Meteorology*, 118, 151–167. [https://doi.org/10.1016/S0168-1923\(03\)00115-1](https://doi.org/10.1016/S0168-1923(03)00115-1)
- Kozłowski, J., & Konarzewski, M. (2005). West, Brown and Enquist's model of allometric scaling again: The same questions remain. *Functional Ecology*, 19, 739–743. <https://doi.org/10.1111/j.1365-2435.2005.01021.x>
- Kozłowski, T. T. (1992). Carbohydrate sources and sinks in woody plants. *Botanical Review*, 58, 107–222. <https://doi.org/10.1007/BF02858600>
- Krinner, G., Viovy, N., de Noblet-Ducoudré, N., Ogée, J., Polcher, J., Friedlingstein, P., ... Prentice, I. C. (2005). A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system. *Global Biogeochemical Cycles*, 19(1), 1–33. <https://doi.org/10.1029/2003GB002199>
- Kuptz, D., Fleischman, F., Matyssek, R., & Grams, T. E. (2011). Seasonal patterns of carbon allocation to respiratory pools in 60-yr-old deciduous (*Fagus sylvatica*) and evergreen (*Picea abies*) trees assessed via whole-tree stable carbon isotope labelling. *New Phytologist*, 191, 160–172. <https://doi.org/10.1111/j.1469-8137.2011.03676.x>
- Li, H. T., Han, X. G., & Wu, J. G. (2005). Lack of evidence for ¾ scaling of metabolism in terrestrial plants. *Journal of Integrative Plant Biology*, 47, 1173–1183. <https://doi.org/10.1111/j.1744-7909.2005.00167.x>
- Litton, C., Raich, J., & Ryan, M. G. (2007). Carbon allocation in forest ecosystems. *Global Change Biology*, 13, 2089–2109. <https://doi.org/10.1111/j.1365-2486.2007.01420.x>
- Lotscher, M., Klumpp, K., & Schnyder, H. (2004). Growth and maintenance respiration for individual plants in hierarchically structured

- canopies of *Medicago sativa* and *Helianthus annuus*: The contribution of current and old assimilates. *New Phytologist*, 164, 305–316.
- Luyssaert, S., Inglima, I., Jung, M., Richardson, A. D., Reichstein, M., Papale, D., ... Janssens, I. A. (2007). CO₂ balance of boreal, temperate, and tropical forests derived from a global database. *Global Change Biology*, 13, 2509–2537.
- Lynch, D. J., Matamala, R., Iversen, C. M., Norby, R. J., & Gonzalez-Meler, M. A. (2013). Stored carbon partly fuels fine-root respiration but is not used for production of new fine roots. *New Phytologist*, 199, 420–430. <https://doi.org/10.1111/nph.12290>
- Machado, J.-L., & Reich, P. B. (2006). Dark respiration rate increases with plant size in saplings of three temperate tree species despite decreasing tissue nitrogen and nonstructural carbohydrates. *Tree Physiology*, 26, 915–923. <https://doi.org/10.1093/treephys/26.7.915>
- Maier, C. A., Johnsen, K. H., Clinton, B. D., & Ludovici, K. H. (2010). Relationships between stem CO₂ efflux, substrate supply, and growth in young loblolly pine trees. *New Phytologist*, 185, 502–513. <https://doi.org/10.1111/j.1469-8137.2009.03063.x>
- Makarieva, A. M., Gorshkov, V. G., & Li, B.-L. (2005). Revising the distributive network models of West, Brown & Enquist (1997) and Banavar, Maritan & Rinaldo (1999). Metabolic inequity of living tissues provides clues for the observed allometric scaling rules. *Journal of Theoretical Biology*, 237, 291–301. <https://doi.org/10.1016/j.jtbi.2005.04.016>
- Makarieva, A. M., Gorshkov, V. G., Li, B.-L., Chown, S. L., Reich, P. B., & Gavrillov, V. M. (2008). Mean mass-specific metabolic rates are strikingly similar across life's major domains: Evidence for life's metabolic optimum. *Proceedings of the National Academy of Sciences of the United States of America*, 105(44), 16994–16999. <https://doi.org/10.1073/pnas.0802148105>
- Mäkelä, A., & Valentine, H. (2001). The ratio of NPP to GPP: Evidence of change over the course of stand development. *Tree Physiology*, 21, 1015–1030. <https://doi.org/10.1093/treephys/21.14.1015>
- Malhi, Y. (2012). The productivity, metabolism and carbon cycle of tropical forest vegetation. *Journal of Ecology*, 100, 65–75. <https://doi.org/10.1111/j.1365-2745.2011.01916.x>
- Malhi, Y., Doughty, C. E., Goldsmith, G. R., Metcalfe, D. B., Girardin, C. A., Marthews, T. R., ... da Costa, A. C. (2015). The linkages between photosynthesis, productivity, growth and biomass in lowland Amazonian forests. *Global Change Biology*, 21, 2283–2295. <https://doi.org/10.1111/gcb.12859>
- Marconi, S., Chiti, T., Nölé, A., Valentini, R., & Collalti, A. (2017). The role of respiration in estimation of net carbon cycle: Coupling soil carbon dynamics and canopy turnover in a novel version of 3D-CMCC forest ecosystem model. *Forests*, 8, 220–227. <https://doi.org/10.3390/f8060220>
- Martínez-Vilalta, J., Sala, A., Asensio, D., Galiano, L., Hoch, G., Palacio, S., ... Lloret, F. (2016). Dynamics of non-structural carbohydrates in terrestrial plants: A global synthesis. *Ecological Monographs*, 86(4), 495–516. <https://doi.org/10.1002/ecm.1231>
- McCree, K. (1970). An equation for the rate of respiration of white clover plants grown under controlled conditions. In I. Šetlík (Ed.), *Prediction and measurement of photosynthetic productivity* (pp. 221–229). Wageningen, The Netherlands: Centre for Agricultural Publishing and Documentation.
- McDowell, N., Pockman, W. T., Allen, C. D., Breshears, D. D., Cobb, N., Kolb, T., ... Yepez, E. A. (2008). Mechanisms of plant survival and mortality during drought: Why do some plants survive while others succumb to drought? *New Phytologist*, 178, 719–739. <https://doi.org/10.1111/j.1469-8137.2008.02436.x>
- McMurtrie, R. E., Norby, R. J., Medlyn, B. E., Dewar, R. C., Pepper, D. A., Reich, P. B., & Barton, C. V. M. (2008). Why is plant-growth response to elevated CO₂ amplified when water is limiting, but reduced when nitrogen is limiting? A growth-optimisation hypothesis. *Functional Plant Biology*, 35, 521–534. <https://doi.org/10.1071/FP08128>
- Medlyn, B., & Dewar, R. (1999). Comment on the article by R. H. Waring, J. J. Landsberg and M. Williams relating net primary production to gross primary production. *Tree Physiology*, 19(2):137–138. <https://doi.org/10.1093/treephys/19.2.137>
- Mencuccini, M., Martínez-Vilalta, J., Vanderklein, D., Hamid, H. A., Korakaki, E., Lee, S., & Michiels, B. (2005). Size-mediated ageing reduces vigour in trees. *Ecology Letters*, 8, 1183–1190. <https://doi.org/10.1111/j.1461-0248.2005.00819.x>
- Merganičová, K., Merganič, J., Lehtonen, A., Vacchiano, G., Zorana, M., Sever, O., ... Reyer, C. P. O. (2019). Forest carbon allocation modelling under climate change. *Tree Physiology*. <https://doi.org/10.1093/treephys/tpz105>
- Michalet, S. T., Cheng, D., Kerkhoff, A. J., & Enquist, B. J. (2014). Convergence of terrestrial plant production across global climate gradients. *Nature*. <https://doi.org/10.1038/nature13470>
- Mori, S., Yamaji, K., Ishida, A., Prokushkin, S. G., Masyagina, O. V., Hagihara, A., ... Umari, M. (2010). Mixed-power scaling of whole-plant respiration from seedlings to giant trees. *Proceedings of the National Academy of Sciences of the United States of America*, 107(4), 1447–1451. <https://doi.org/10.1073/pnas.0902554107>
- Muhr, J., Angert, A., Negron-Juarez, R. I., Munoz, W. A., Kraemer, G., Chambers, J. Q., & Trumbore, S. E. (2013). Carbon dioxide emitted from live stems of tropical trees is several years old. *Tree Physiology*, 33, 743–752. <https://doi.org/10.1093/treephys/tpt049>
- Muhr, J., Messier, C., Delagrangé, S., Trumbore, S., Xu, X., & Hartmann, H. (2016). How fresh is maple syrup? Sugar maple trees mobilize carbon stored several years previously during early springtime sap-ascend. *New Phytologist*, 209, 1410–1416. <https://doi.org/10.1111/nph.13782>
- O'Connor, M. P., Kemp, S. J., Agosta, S. J., Hansen, F., Sieg, A. E., Wallace, B. P., ... Dunham, A. E. (2007). Reconsidering the mechanistic basis of the metabolic theory of ecology. *Oikos*, 116, 1058–1072. <https://doi.org/10.1111/j.2006.0030-1299.15534.x>
- Odum, E. P. (1969). The strategy of ecosystem development. *Science*, 164(3877), 262–270. <https://doi.org/10.1126/science.164.3877.262>
- Oleson, K. W., Lawrence, D. M., Bonan, G. B., Drewniak, B., Huang, M., Koven, C. D., ... Yang, Z.-L. (2013). *Technical description of version 4.5 of the community land model (CLM)*. Ncar Technical Note NCAR/TN-503+STR. Boulder, CO: National Center for Atmospheric Research, 422 pp.
- Palacio, S., Hernández, R., Maestro-Martínez, M., & Camarero, J. J. (2012). Fast replenishment of initial carbon stores after defoliation by the pine processionary moth and its relationship to the re-growth ability of trees. *Trees – Structure and Function*, 26, 1627–1640. <https://doi.org/10.1007/s00468-012-0739-y>
- Pallardy, S. G. (2010). *Physiology of woody plants*. New York, NY: Academic Press.
- Pappas, C., Fatichi, S., Leuzinger, S., Wolf, A., & Burlando, P. (2013). Sensitivity analysis of a process-based ecosystem model: Pinpointing parameterization and structural issues. *Journal of Geophysical Research*, 118, 502–528. <https://doi.org/10.1002/jgrg.20035>
- Penning de Vries, F. W. T. (1975). The cost of maintenance processes in plants. *Annals of Botany*, 39, 77–92. <https://doi.org/10.1093/oxfordjournals.aob.a084919>
- Penning de Vries, F. W. T., Brunsting, A. H. M., & van Laar, H. H. (1974). Products, requirements and efficiency of biosynthesis: A quantitative approach. *Journal of Theoretical Biology*, 45, 339–377.
- Piao, S., Luyssaert, S., Ciais, P., Janssens, I. A., Chen, A., Cao, C., ... Wang, S. (2010). Forest annual carbon cost: A global-scale analysis of autotrophic respiration. *Ecology*, 91(3), 652–661. <https://doi.org/10.1890/08-2176.1>
- Pietsch, S. A., Hasenauer, H., & Thornton, P. E. (2005). BGC-model parameters for tree species growing in central European forests. *Forest*

- Ecology and Management*, 211, 264–295. <https://doi.org/10.1016/j.foreco.2005.02.046>
- Poulter, B., Hattermann, F., Hawkins, E. D., Zaehle, S., Sitch, S., Restrepo-Coupe, N. A., ... Cramer, W. (2010). Robust dynamics of Amazon dieback to climate change with perturbed ecosystem model parameters. *Global Change Biology*, 16, 2476–2495. <https://doi.org/10.1111/j.1365-2486.2009.02157.x>
- Price, C. A., Gilooley, J. F., Allen, A. P., Weitz, J. S., & Niklas, K. J. (2010). The metabolic theory of ecology: Prospects and challenges for plant biology. *New Phytologist*, 188, 696–710. <https://doi.org/10.1111/j.1469-8137.2010.03442.x>
- Price, C. A., Wietz, J. S., Savage, V. M., Stegen, J., Clarke, A., Coomes, D. A., ... Niklas, K. J. (2012). Testing the metabolic theory of ecology. *Ecology Letters*, 15, 1465–1474. <https://doi.org/10.1111/j.1461-0248.2012.01860.x>
- Pringle, E. G. (2016). Integrating plant carbon dynamics with mutualism ecology. *New Phytologist*, 210, 71–75.
- Reich, P. B., Tjoelker, M. G., Machado, J.-L., & Oleksyn, J. (2006). Universal scaling of respiratory metabolism, size and nitrogen in plants. *Nature*, 439, 457–461. <https://doi.org/10.1038/nature04282>
- Reich, P. B., Tjoelker, M. G., Pregitzer, K. S., Wright, I. J., Oleksyn, J., & Machado, J.-L. (2008). Scaling of respiration to nitrogen in leaves, stems and roots of higher land plants. *Ecology Letters*, <https://doi.org/10.1111/j.1461-0248.2008.01185.x>
- Reyer, C., Silveyra, G. R., Dolos, K., Hartig, F., Hauf, Y., & Noack, M., ... Frieler, K. (2019). The PROFOUND database for evaluating vegetation models and simulating climate impacts on forests. V. 0.1.12. GFZ Data Services. <https://doi.org/10.5880/PIK.2019.008>
- Richardson, A. D., Carbone, M. S., Hugggett, B. A., Furze, M. E., Czimeczik, C. I., Walker, J. C., ... Murakami, P. (2015). Distribution and mixing of old and new nonstructural carbon in two temperate trees. *New Phytologist*, 206, 590–597. <https://doi.org/10.1111/nph.13273>
- Richardson, A. D., Carbone, M. S., Keenan, T. F., Czimeczik, C. I., Hollinger, D. Y., Murakami, P., ... Xu, X. (2013). Seasonal dynamics and age of stemwood nonstructural carbohydrates in temperate forest trees. *New Phytologist*, 197, 850–861. <https://doi.org/10.1111/nph.12042>
- Roxburgh, S. H., Berry, S. L., Buckley, T. N., Barnes, B., & Roderick, M. L. (2005). What is NPP? Inconsistent accounting of respiratory fluxes in the definition of net primary production. *Functional Ecology*, 19, 378–382. <https://doi.org/10.1111/j.1365-2435.2005.00983.x>
- Sala, A., Fouts, W., & Hoch, G. (2011). Carbon storage in trees: Does relative carbon supply decrease with tree size? In F. C. Meinzer, B. Lachenbruch, & T. E. Dawson (Eds.), *Size- and age-related changes in tree structure and function* (pp. 287–306). Dordrecht, the Netherlands: Springer.
- Sala, A., Woodruff, D. R., & Meinzer, F. C. (2012). Carbon dynamics in trees: Feast or famine? *Tree Physiology*, 32, 764–775. <https://doi.org/10.1093/treephys/tp143>
- Sato, H., Itoh, A., & Kohyama, T. (2007). SEIB-DGVM: A new Dynamic Global Vegetation Model using a spatially explicit individual-based approach. *Ecological Modelling*, 200, 279–307. <https://doi.org/10.1016/j.ecolmodel.2006.09.006>
- Schiestl-Aalto, P., Kulmala, L., Nikinmaa, E., & Mäkelä, A. (2015). CASSIA – A dynamic model for predicting intra-annual sink demand and interannual growth variation in Scots pine. *New Phytologist*, 206, 647–659. <https://doi.org/10.1111/nph.13275>
- Schiestl-Aalto, P., Ryhti, K., Mäkelä, A., Peltoniemi, M., Bäck, J., & Kulmala, L. (2019). Analysis of the NSC storage dynamics in tree organs reveals the allocation to belowground symbionts in the framework of whole tree carbon balance. *Frontiers in Forests and Global Change*, 2, 17. <https://doi.org/10.3389/ffgc.2019.00017>
- Schuur, E. A. G., & Trumbore, S. E. (2006). Partitioning sources of soil respiration in boreal black spruce forest using radiocarbon. *Global Change Biology*, 12, 165–176. <https://doi.org/10.1111/j.1365-2486.2005.01066.x>
- Schwalm, C. R., & Ek, A. R. (2004). A process-based model of forest ecosystems driven by meteorology. *Ecological Modelling*, 179, 317–348. <https://doi.org/10.1016/j.ecolmodel.2004.04.016>
- Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., ... Thonicke, K. (2003). Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model. *Global Change Biology*, 9, 161–185. <https://doi.org/10.1046/j.1365-2486.2003.00569.x>
- Smith, A. M., & Stitt, M. (2007). Coordination of carbon supply and plant growth. *Plant, Cell & Environment*, 30, 1126–1149. <https://doi.org/10.1111/j.1365-3040.2007.01708.x>
- Smith, B., Prentice, I. C., & Sykes, M. T. (2001). Representation of vegetation dynamics in the modelling of terrestrial ecosystems: Comparing two contrasting approaches within European climate space. *Global Ecology and Biogeography*, 10, 621–637. <https://doi.org/10.1046/j.1466-822X.2001.00256.x>
- Solly, E. F., Brunner, I., Helmisaari, H. S., Herzog, C., Leppälammil-Kujansuu, J., Schöning, I., ... Hagedorn, F. (2018). Unravelling the age of fine roots of temperate and boreal forests. *Nature Communications*, 9.
- Spicer, R., & Holbrook, N. M. (2007). Parenchyma cell respiration and survival in secondary xylem: Does metabolic activity decline with cell age? *Plant, Cell & Environment*, 30, 934–943. <https://doi.org/10.1111/j.1365-3040.2007.01677.x>
- Stephenson, N. L., Das, A. J., Condit, R., Russo, S. E., Baker, P. J., Beckman, N. G., ... Alvarez, E. (2014). Rate of tree carbon accumulation increases continuously with tree size. *Nature*, 507, 90–93. <https://doi.org/10.1038/nature12914>
- Tang, J., Luyssaert, S., Richardson, A. D., Kutsch, W., & Janssens, I. A. (2014). Steeper declines in forest photosynthesis than respiration explain age-driven decreases in forest growth. *Proceedings of the National Academy of Sciences of the United States of America*, 111(24), 8856–8860. <https://doi.org/10.1073/pnas.1320761111>
- Tatarinov, F., & Cenciala, E. (2006). Application of BIOME-BGC model to managed forests 1. Sensitivity analysis. *Forest Ecology and Management*, 237, 267–279. <https://doi.org/10.1016/j.foreco.2006.09.085>
- Thornley, J. H. M. (1970). Respiration, growth and maintenance in plants. *Nature*, 227, 304–305. <https://doi.org/10.1038/227304b0>
- Thornley, J. H. M. (2011). Plant growth and respiration re-visited: Maintenance respiration defined – It is an emergent property of, not a separate process within, the system – And why the respiration: Photosynthesis ratio is conservative. *Annals of Botany*, 108, 1365–1380. <https://doi.org/10.1093/aob/mcr238>
- Thornley, J. H. M., & Cannell, M. G. R. (2000). Modelling the components of plant respiration: Representation and realism. *Annals of Botany*, 85, 55–67. <https://doi.org/10.1006/anbo.1999.0997>
- Thornton, P. E., Law, B. E., Gholz, H. L., Clark, K. L., Falge, E., Ellsworth, D. S., ... Chen, J. (2002). Modeling and measuring the effects of disturbance history and climate on carbon and water budgets in evergreen needleleaf forests. *Agricultural and Forest Meteorology*, 113, 185–222. [https://doi.org/10.1016/S0168-1923\(02\)00108-9](https://doi.org/10.1016/S0168-1923(02)00108-9)
- Turner, M., Beer, C., Ciais, P., Friend, A. D., Ito, A., Kleidon, A., ... Carvalhais, N. (2017). Evaluation of climate-related carbon turnover processes in global vegetation models for boreal and temperate forests. *Global Change Biology*, 23(8), 3076–3091. <https://doi.org/10.1111/gcb.13660>
- Tjoelker, M. G., Oleksyn, J., & Reich, P. B. (1999). Acclimation of respiration to temperature and CO₂ in seedlings of boreal tree species in relation to plant size and relative growth rate. *Global Change Biology*, 49, 679–691. <https://doi.org/10.1046/j.1365-2486.1999.00257.x>
- Van Oijen, M., Schapendonk, A., & Hoglind, M. (2010). On the relative magnitudes of photosynthesis, respiration, growth and carbon storage in vegetation. *Annals of Botany*, 105, 739–797. <https://doi.org/10.1093/aob/mcq039>

- Vargas, R., Trumbore, S. E., & Allen, M. F. (2009). Evidence of old carbon used to grow new fine roots in a tropical forest. *New Phytologist*, 182, 710–718. <https://doi.org/10.1111/j.1469-8137.2009.02789.x>
- Waring, R. H., Landsberg, J. L., & Williams, M. (1998). Net primary production of forests: A constant fraction of gross primary production? *Tree Physiology*, 18, 129–134. <https://doi.org/10.1093/treephys/18.2.129>
- Warnant, P., François, L., Strivay, D., & Gérard, J.-C. (1994). CARAIB: A global model of terrestrial biological productivity. *Global Biogeochemical Cycles*, 8(3), 255–270. <https://doi.org/10.1029/94gb00850>
- Warszawski, L., Frieler, K., Huber, V., Piontek, F., Serdeczny, O., & Schewe, J. (2014). The Inter-Sectoral Impact Model Intercomparison Project (ISI-MIP): Project framework. *Proceedings of the National Academy of Sciences of the United States of America*, 111(9), 3228–3232. <https://doi.org/10.1073/pnas.1312330110>
- Weber, R., Schwendener, A., Schmid, S., Lambert, S., Wiley, E., Landhäusser, S. M., ... Hoch, G. (2018). Living on next to nothing: Tree seedlings can survive weeks with very low carbohydrate concentration. *New Phytologist*, 218, 107–118. <https://doi.org/10.1111/nph.14987>
- West, G. B., Brown, J. H., & Enquist, B. J. (1999). A general model for the structure and allometry of plant vascular systems. *Nature*, 400, 664–667. <https://doi.org/10.1038/23251>
- White, M., Thornton, P. E., Running, S. W., & Nemani, R. R. (2000). Parameterization and sensitivity analysis of the BIOME-BGC terrestrial ecosystem model: Net primary production controls. *Earth Interactions*, 4, 1–85. [https://doi.org/10.1175/1087-3562\(2000\)004<0003:pasaot>2.0.co;2](https://doi.org/10.1175/1087-3562(2000)004<0003:pasaot>2.0.co;2)
- Wiley, E., & Helliker, B. (2012). A re-evaluation of carbon storage in trees lends greater support for carbon limitation to growth. *New Phytologist*, 195, 285–289. <https://doi.org/10.1111/j.1469-8137.2012.04180.x>
- Wiley, E., Hoch, G., & Landhäusser, S. M. (2017). Dying piece by piece: Carbohydrate dynamics in aspen (*Populus tremuloides*) seedlings under severe carbon stress. *Journal of Experimental Botany*, 68(18), 5221–5232. <https://doi.org/10.1093/jxb/erx342>
- Woodruff, D. R., & Meinzer, F. C. (2011). Water stress, shoot growth and storage of nonstructural carbohydrates along a tree height gradient in a tall conifer. *Plant, Cell & Environment*, 11, 1920–1930. <https://doi.org/10.1111/j.1365-3040.2011.02388.x>
- Wu, J., Larsen, K., van der Linden, L., Beier, C., Pilegaard, K., & Ibrom, A. (2013). Synthesis on the carbon budget and cycling in a Danish, temperate deciduous forest. *Agricultural and Forest Meteorology*, 181, 94–107. <https://doi.org/10.1016/j.agrformet.2013.07.012>
- Wythers, K. R., Reich, P. B., & Bradford, J. B. (2013). Incorporating temperature-sensitive Q10 and foliar respiration acclimation algorithms modify modeled ecosystem responses to global change. *Journal of Geophysical Research: Biogeosciences*, 118, 1–14. <https://doi.org/doi:10.1029/2011JG001897>
- Xu, C.-Y., Turnbull, M. H., Tissue, D. T., Lewis, J. D., Carson, R., Schuster, W. S. F., ... Griffin, K. L. (2012). Age-related decline of stand biomass accumulation is primarily due to mortality and not to reduction in NPP associated with individual tree physiology, tree growth or stand structure in a *Quercus*-dominated forest. *Journal of Ecology*, 100, 428–440. <https://doi.org/10.1111/j.1365-2745.2011.01933.x>
- Yang, J., He, Y., Aubrey, D. P., Zhuang, Q., & Teskey, R. O. (2016). Global patterns and predictors of stem CO₂ efflux in forest ecosystems. *Global Change Biology*, 22, 1433–1444. <https://doi.org/10.1111/gcb.13188>
- Yang, Y., Luo, Y., & Finzi, A. C. (2011). Carbon and nitrogen dynamics during forest stand development: A global synthesis. *New Phytologist*, 190, 977–989. <https://doi.org/10.1111/j.1469-8137.2011.03645.x>
- Zaehle, S., Sitch, S., Smith, B., & Hatterman, F. (2005). Effects of parameter uncertainties on the modelling of terrestrial biosphere dynamics. *Global Biogeochemical Cycles*, 19, 1–16. <https://doi.org/10.1029/2004GB002395>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Collalti A, Tjoelker MG, Hoch G, et al. Plant respiration: Controlled by photosynthesis or biomass? *Glob Change Biol*. 2020;26:1739–1753. <https://doi.org/10.1111/gcb.14857>